## BODY CONTOURING SURGERY:

# Brachioplasty (Arm Lift), Thigh Lift & Buttock Lift

### What to Expect

#### **OPERATION DURATION**

This operation is performed under general anaesthetic and usually takes 3-4 hours (per bilateral body area.)

#### HOSPITAL STAY

You will be in the hospital for 1 - 2 nights, depending on your recovery. Everyone is different, and your recovery will be tailored to you.

Dr Taylor will not be in to visit you every day whilst you are in the hospital. He will phone your nurses for daily updates and provide over-the-phone instructions. He will come in to see you if you have any issues.

Our specialist nurse will follow up with you on the phone once you have been discharged. You will also be given her direct contact number. Feel free to phone or message her 9 am - 5 pm, 7 days a week, if you have any questions or concerns.

### What Complications May Occur After This Procedure?

## GENERAL POSSIBLE COMPLICATIONS OF ANY PROCEDURE

- Bleeding
- Infection at the surgical site
- Allergic reaction to equipment, materials or medication
- Blood clot in your leg or lungs
- Acute kidney injury
- Poor scarring hypertrophic or keloid

## SPECIFIC POSSIBLE COMPLICATIONS OF THIS PROCEDURE

- Lump developing under your wound caused by blood collecting (haematoma) that may require further surgery
- Excess fluid accumulating under the skin (seroma) may require draining
- Slow-healing wounds or areas of fat necrosis that may require regular dressings for an extended period
- Skin necrosis, where some of the skin at the edge of your wound dies and leaves a black area
- Wound healing problems wound has delayed healing and may open up (dehiscence)
- Numbness or continued pain around the scars
- Restrictive movement to the tightened skin
- Asymmetry

## SPECIFIC POSSIBLE COMPLICATIONS OF THIS PROCEDURE

#### Thigh Lift Only

• Distortion of the genitalia's contour and appearance, especially in women

#### **Buttock Lift Only**

- Irregular skin surface, uneven contours or rippling
- Asymmetry (unevenness) of the buttocks

#### CONSEQUENCES OF THIS PROCEDURE

- Pain, bruising and swelling
- Visible scars

## How to Prepare for Your Procedure

#### HOW LONG SHOULD YOU BOOK OFF WORK

This depends on your profession - If you work in an office/ non-strenuous job 1-2 weeks. However, this will depend on how physically demanding your job is, so please discuss it with Dr Taylor or our nurse.

Please email <u>admin@drduncantaylor.com.au</u> if you require a medical certificate.

#### SUPPORT GARMENT

You will need to purchase a post-surgery compression garment:

Brachioplasty - Arm garments

Buttock or Thigh Lift - Knee-length compression shorts

Please bring these into the theatre with you on the day of surgery.

Wear your garments 24/7 for 6 weeks (including in the hospital) you can remove the garments to shower.

Breast Care WA After Surgery Garment & Fitting Specialists (<u>www.breastcarewa.com.au</u>)

#### FASTING

You are required to fast for General Anaesthesia. You cannot have any food or fluids (including lollies, chewing gum) 6 hours prior to your surgery. You can have some sips of water only up to 2 hours before your surgery and then nothing after this time.

If you take regular medication, please check with our nurse to see if you should be taking your medication in the lead-up to/on the day of your surgery. Some medications should be withheld.

#### PRE-OP WASH

Please purchase a pre-op chlorhexidine wash from the pharmacy. Use this whilst showering, the night before and the morning of your surgery.

#### MORNING OF SURGERY

You will be asked to arrive at the hospital approximately 2 hours prior to your surgery start time. Our Practice Manager will email you your admission information, including time and where to go, 1-2 weeks before your surgery date.

Dr Taylor will see you in the pre-operative area before you go into theatre. He will mark out the site for your surgery. This process must be completed whilst you are awake, as he requires you to be standing.

#### DRAINS

You may have drains inserted into your wounds immediately post-op. If this is the case, these will be removed before your hospital discharge.

## **Post-Operative Instructions**

#### ACTIVITY

You can get up and move the same day after your surgery once you have recovered from your general anaesthetic.

In the initial 2 weeks following your surgery, we would like you to rest the area of your body as much as possible to assist with your wound healing. Pottering around the house is fine, though do not start going for walks, doing housework, etc. during this time.

After 2 weeks, you can start some short/slow walks and slowly increase from 2 - 6 weeks post-op.

Please do not do any strenuous physical activity or heavy lifting over 10 kg for 3 weeks. Walking is fine. You cannot drive for 24 hours after your general anaesthetic.

#### PAIN RELIEF

You should not be in significant pain after your surgery. Your anaesthetist will prescribe regular pain relief for on the ward post-surgery. If you have pain on top of this, you should ask for additional pain medication, which will be prescribed on your drug chart should you need it.

You will not be discharged from the hospital until your pain is well managed on oral medications.

Please take paracetamol regularly and your stronger pain relief, if required. If you have severe pain that is not responding to the pain relief, please phone our nurse during office hours.

Please do not put ice packs on your wounds.

#### Do not take aspirin.

#### ANTI-INFLAMMATORIES

You will be prescribed some anti-inflammatories whilst you are in hospital, and you will also be discharged with some. Please take the entire course, even if you are not in pain. Please take this medication as prescribed on the box by your anaesthetist.

#### **ANTIBIOTICS**

You will be given a 5 days course of oral antibiotics. Depending on how long you remain in the hospital,

you may be discharged home with some remaining antibiotics. Please take the full course.

#### BLEEDING

If the wound starts to bleed, please apply firm pressure for 15 minutes. If a dressing is in place, please do this over the top of the dressing (do not remove the dressing).

This should control the bleeding. However, if it is not easily controlled or does not subside, please phone our nurse specialist during working hours, go to an afterhours GP clinic or visit an emergency department.

#### **INFECTION**

If your wound becomes hot, swollen, oozes pus, or you are spiking temperatures, the wound may be infected. Please phone our nurse immediately within business hours or attend your GP, after-hours GP or Emergency.

#### **REMOVAL OF SUTURES**

Sutures used during these procedures do not require removing and they will dissolve on their own.

#### WOUND & DRESSING CARE

Your wounds will be covered with surgical tape and glue, this is waterproof, and you can shower as usual and pat dry. This dressing will start to peel off on its own after approximately 3 weeks. At this stage, you can gently peel the dressing off in the shower.

If you have a standard dressing on, please sponge bath around these to avoid getting the dressing wet and leave intact until you are seen in the nurse dressing clinic 5-10 days post-op.

Do not immerse the wound in water, i.e.. swimming or bathing for 6 weeks, or until the wound is completely healed.

Bruising takes around 2-3 weeks to resolve, and swelling can take up to 3-6 months to resolve completely.

#### MASSAGING FOR SCAR REDUCTION

Commence massaging your scars once your dressings are removed, and wounds have healed – approximately 3 weeks post-op.

It is recommended that you massage your scars with emollient or bio oil for 2 minutes, 2-3 times per day and then apply a silicone gel to sit directly on the wound (this sits on the scar and does not get massaged in). Continue this for 3-6 months.

If you would like assistance with scar management and massage therapy, we can refer you to an occupational therapist. This service does incur an out-of-pocket expense and is not necessary for all patients, though if this is something you are interested in, please inform our nurse.

## Other Useful Tips

The power outlet in your hospital room is often quite far away from your bed. We suggest bringing a long charging cable for your phone so that you can keep it within reach.

## Follow-Up

Dr Taylor will be in contact with the nurses looking after you on the ward. If you are well post-op he will discharge you after speaking with your nurse over the phone. He will only come in to see you on the ward if you are having any issues. Our nurse will phone you a day or 2 after your discharge to check in on you (this call may be after hours).

If you are concerned at all or have any questions, please phone our clinic to speak with our practice nurse.

#### Follow up appointments:

- 5-10 days after your procedure, post-surgery wound check in our nurse clinic.
- 6 weeks post-surgery review in Dr Duncan Taylor's day clinic.

#### If you have any questions please contact us:

Phone (08) 9439 7406 | Fax 08 6266 3751
Email admin@drduncantaylor.com.au
Healthlink DMTaylor
Rooms 106 Outram Street, West Perth, WA, 6005

For more information visit <u>drduncantaylor.com.au</u>









/ Australian Society of Plastic Surgeons

